

# EFFECTS OF ORGANIC MATTER ON THE SPECIATION OF URANIUM IN SOIL AND PLANT MATRICES

A.J. Bednar\*, V.F. Medina, and S.L. Larson  
US Army Engineer Research and Development Center  
Vicksburg, MS 39180

D.S. Ulmer-Scholle and B.A. Frey  
New Mexico Institute of Mining and Technology  
Socorro, NM 87801

J.G. Moran  
New Mexico State University  
Las Cruces, NM 88003

## ABSTRACT

Radionuclides are known to complex with organic matter, which can promote mobility in soil environments. This work focuses on interactions of depleted uranium with organic compounds using HPLC-ICP-MS to identify organouranium species in soil and plant materials. Nearly all of the uranium extracted from certain plant tissues is bound to organic ligands. These experiments suggest organic compounds may be a significant influence on the chemistry of uranium in the environment.

## 1. INTRODUCTION

Metal complexation with ligands has been known to affect physical and chemical properties of the affected metal (Nierop, et al., 2002). Natural organic matter (NOM), for example, contains functional groups that can form complexes with metals (Christensen et al., 1999; Chen et al., 2003). Radionuclides, in particular, have been studied and their interactions with NOM reported in the literature (Zhang et al., 1997; Lenhart et al., 2000, Pacheco and Havel, 2001).

Anthropogenic activities, such as mining, milling, and enrichment for nuclear fuel, can increase the potential mobility of uranium (Abdelouas et al., 1998). Additionally, the waste material from the enrichment process, isotopically 'depleted' uranium (DU), also finds uses, particularly in armor and armor piercing munitions (Chen and Yiacoumi, 2002; Sztajnkrzyer and Otten, 2004). The United States Army does not use DU munitions for training, but does conduct testing, and DU has been historically used in military conflicts, which can result in the release of metallic DU into the environment (Sztajnkrzyer and Otten, 2004). Although metallic uranium is essentially immobile, corrosion reactions can yield oxidized products (Chen and Yiacoumi, 2002), which can dissolve and complex with NOM. This work

investigates the extent of organouranium complexes in various plant and soil matrices.

## 2. EXPERIMENTAL

An Agilent series 1100 High Pressure Liquid Chromatograph (HPLC) was used for all separations and was coupled to a Perkin Elmer Elan 6000 ICP-MS. Reagent grade chemicals and 18.2 M $\Omega$ cm deionized water were used for preparation of mobile phases, and all solutions were filtered (0.45  $\mu$ m) prior to introduction into the instrument.

Molecular size exclusion (SE), or gel permeation, chromatography is an on-line size separation technique. A Shodex KW-803 SEC column was used to separate organouranium complexes based on molecular weight (hydrodynamic size). The mobile phase used in all experiments was 21 mM Tris buffer, (pH = 7) because of the protein molecular weight standards used to correlate elution time with molecular size. An ultraviolet-visible (UV) absorbance detector was added in-line prior to the ICP-MS to detect organic moieties. This dual detector capability allowed simultaneous determination of organic moieties that contain uranium and those that do not.

## 3. RESULTS AND DISCUSSION

Fig. 1 is a size exclusion chromatogram simultaneously using ICP-MS as an elemental specific detector for uranium and UV absorbance (230 nm), as a surrogate measurement for organic matter (SEC-UV-ICP-MS). The plant material in Fig. 1 is Indian Mustard from a phytoremediation study. Both inorganic and organouranium forms are seen in this plant material. The SEC-UV-ICP-MS chromatogram shown in Fig. 2 is of a cryptobiotic soil crust deionized water extract. These samples show the complex nature of the organouranium species extracted by deionized water. The soil crust sample contains a high level of uranium (~100  $\mu$ g/L) with multiple inorganic and organouranium species. The

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>00 DEC 2004</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Effects Of Organic Matter On The Speciation Of Uranium In Soil And Plant Matrices</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>US Army Engineer Research and Development Center Vicksburg, MS 39180; New Mexico Institute of Mining and Technology Socorro, NM 87801</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM001736, Proceedings for the Army Science Conference (24th) Held on 29 November - 2 December 2005 in Orlando, Florida.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>UU</b>	18. NUMBER OF PAGES <b>2</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

uranium peak observed at ~6 min appears to be an inorganic colloidal form, as it did not have a significant UV signature, whereas the peaks observed at ~10-15 min consisted of uranium associated with the organic moieties.




Fig. 1: Size exclusion chromatogram of organouranium complexes extracted from Indian Mustard plant material.




Fig. 2: Size exclusion chromatogram of organouranium complexes extracted from soil crust material.

The most notable information gained from the SEC data in Figs. 1 and 2 was the distribution of uranium in these samples. Large molecules elute earlier than smaller ones as they are 'excluded' from pore spaces in the column packing. The data for the plant extract (Fig. 1) and the soil crust extract (Fig. 2) indicate that the inorganic uranium peaks (those without a large UV absorbance) were in particulate form and of a larger size than the organouranium complexes (because they elute first at about 6 minutes). Analysis of protein MW standards on this column with this mobile phase suggest the organouranium complexes eluting at 10-15 minutes have similar hydrodynamic size to ~ 3 kDa MW proteins.

## ACKNOWLEDGEMENTS

The use of trade or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government. The tests described and the data presented herein, were obtained from research conducted under the Environmental Quality Technology Program of the USACE by the USAERDC. Permission was granted by the Chief of Engineers to publish this information. The authors thank Mr. Donald Brown and Mr. Charles White of the USACE for editorial comments.

## REFERENCES

- Abdelouas, A., Lutze, W., and Nuttall, E., 1998: Chemical reactions of uranium in ground water at a mill tailingsite. *J. of Cont. Hydro.*, 34, 343-361.
- Chen, J.P., and Yiacoumi, S., 2002: Modeling of depleted uranium transport in subsurface systems. *Wat. Air and Soil Poll.*, 140, 173-201.
- Christensen, J.B., Botma, J.J., and Christensen, T.H., 1999: Complexation of Cu and Pb by DOC in polluted groundwater: a comparison of experimental data and predictions by computer speciation models (WHAM and MINTEQA2). *Wat. Res.*, 33, 3231-3238.
- Lenhart, J.J., Cabaniss, S.E., MacCarthy, P., and Honeyman, B.D., 2000: Uranium(VI) complexation with citric, humic, and fulvic acids. *Radiochim. Acta.*, 88, 345-353.
- Nierop, K.G.J., Jansen, B., and Verstraten, J.M., 2002: Dissolved organic matter, aluminum, and iron interactions: precipitation induced by metal/carbon ratio, pH, and competition. *Sci. of the Total Envir.*, 300, 201-211.
- Pacheco, M.L., and Havel, J., 2001: Capillary zone electrophoretic (CZE) study of uranium(VI) complexation with humic acids. *J. of Radio. and Nuc. Chem.*, 248, 565-570.
- Sztajnkrzyer, M.D., and Otten, E.J., 2004: Chemical and radiological toxicity of depleted uranium. *Mil. Med.*, 169, 212-216.
- Zhang, Y.J., Bryan, N.D., Livens, F.R., and Jones, M.N., 1997: Selectivity in the complexation of actinides by humic substances. *Envir. Poll.*, 96, 361-367.

## CONCLUSIONS

The experiments presented demonstrate that NOM interactions are important to the chemistry of uranium in natural systems. Size exclusion chromatography was used to investigate uranium binding with organic moieties in plant and soil extracts, and showed multiple inorganic and organouranium forms in natural samples. The results of this work indicate SEC is useful for studying binding of uranium to complex organic moieties found in nature.